

Wintenberger's Functor for Abelian Extensions

Kevin Keating
 Department of Mathematics
 University of Florida
 Gainesville, FL 32611
 USA
 keating@math.ufl.edu

May 20, 2008

Abstract

Let k be a finite field. Wintenberger used the field of norms to give an equivalence between a category whose objects are totally ramified abelian p -adic Lie extensions E/F , where F is a local field with residue field k , and a category whose objects are pairs (K, A) , where $K \cong k((T))$ and A is an abelian p -adic Lie subgroup of $\text{Aut}_k(K)$. In this paper we extend this equivalence to allow $\text{Gal}(E/F)$ and A to be arbitrary abelian pro- p groups.

1 Introduction

Let $q = p^f$ and let k be a finite field with q elements. We define a category $\mathcal{A}$ whose objects are totally ramified abelian extensions E/F , where F is a local field with residue field k , and $[E : F]$ is infinite if F has characteristic 0. An $\mathcal{A}$ -morphism from E/F to E'/F' is defined to be a continuous embedding $\rho : E \rightarrow E'$ such that

1. ρ induces the identity on k .
2. E' is a finite separable extension of $\rho(E)$.
3. F' is a finite separable extension of $\rho(F)$.

Let $\rho^* : \text{Gal}(E'/F') \rightarrow \text{Gal}(E/F)$ be the map induced by ρ . It follows from conditions 2 and 3 that ρ^* has finite kernel and finite cokernel.

We also define a category $\mathcal{B}$ whose objects are pairs (K, A) , where K is a local field of characteristic p with residue field k and A is a closed abelian subgroup of $\text{Aut}_k(K)$. A $\mathcal{B}$ -morphism from (K, A) to (K', A') is defined to be a continuous embedding $\sigma : K \rightarrow K'$ such that

1. σ induces the identity on k .
2. K' is a finite separable extension of $\sigma(K)$.
3. A' stabilizes $\sigma(K)$, and the image of A' in $\text{Aut}_k(K)$ is an open subgroup of A .

Let $\sigma^* : A' \rightarrow A$ be the map induced by σ . It follows from conditions 2 and 3 that σ^* has finite kernel and finite cokernel.

The field of norms construction [7] gives a functor $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$ defined by

$$\mathcal{F}(E/F) = (X_F(E), \text{Gal}(E/F)). \quad (1.1)$$

We wish to prove the following:

Theorem 1.1 *$\mathcal{F}$ is an equivalence of categories.*

Wintenberger ([5, 6]; see also [2]) has shown that $\mathcal{F}$ induces an equivalence between the full subcategory $\mathcal{A}_L$ of $\mathcal{A}$ consisting of extensions E/F such that $\text{Gal}(E/F)$ is an abelian p -adic Lie group, and the full subcategory $\mathcal{B}_L$ of $\mathcal{B}$ consisting of pairs (K, A) such that A is an abelian p -adic Lie group. (Contrary to [2, 6] we consider finite groups to be p -adic Lie groups. The equivalence of categories proved in [2, 5, 6] extends trivially to include the case of finite groups.) The proof of Theorem 1.1 is based on reducing to the equivalence between $\mathcal{A}_L$ and $\mathcal{B}_L$.

The following result, proved by Laubie [3], is a consequence of Theorem 1.1:

Corollary 1.2 *Let $(K, A) \in \mathcal{B}$. Then there is $E/F \in \mathcal{A}$ such that A is isomorphic to $G = \text{Gal}(E/F)$ as a filtered group. That is, there exists an isomorphism $i : A \rightarrow G$ such that $i(A[x]) = G[x]$ for all $x \geq 0$, where $A[x]$, $G[x]$ denote the ramification subgroups of A , G with respect to the lower numbering.*

The finite field $k \cong \mathbb{F}_q$ is fixed throughout the paper, as is the field $K = k((T))$ of formal Laurent series over k . We work with complete discretely valued fields F whose residue field is identified with k , and with totally ramified abelian extensions of such fields. The ring of integers of F is denoted by $\mathcal{O}_F$ and the maximal ideal of $\mathcal{O}_F$ is denoted by $\mathcal{M}_F$. We let v_F denote the valuation on the separable closure F^{sep} of F which is normalized so that $v_F(F^\times) = \mathbb{Z}$, and we let v_p denote the p -adic valuation on $\mathbb{Z}$. We say that the profinite group G is finitely generated if there is a finite set $S \subset G$ such that $\langle S \rangle$ is dense in G .

2 Ramification theory and the field of norms

In this section we recall some facts from ramification theory, and summarize the construction of the field of norms for extensions in $\mathcal{A}$.

Let $E/F \in \mathcal{A}$. Then $G = \text{Gal}(E/F)$ has a decreasing filtration by the upper ramification subgroups $G(x)$, defined for nonnegative real x . (See for instance [4, IV].)

We say that u is an upper ramification break of G if $G(u + \epsilon) \subsetneq G(u)$ for every $\epsilon > 0$. Since G is abelian, by the Hasse-Arf Theorem [4, V §7, Th. 1] every upper ramification break of G is an integer. In addition, since E/F is a totally ramified abelian extension, it follows from class field theory that E/F is arithmetically profinite (APF) in the sense of [7, §1]. This means that for every $x \geq 0$ the upper ramification subgroup $G(x)$ has finite index in $G = G(0)$. This allows us to define the Hasse-Herbrand functions

$$\psi_{E/F}(x) = \int_0^x |G(0) : G(t)| dt \quad (2.1)$$

and $\phi_{E/F}(x) = \psi_{E/F}^{-1}(x)$. It follows that the ramification subgroups of G with the lower numbering can be defined by $G[x] = G(\phi_{E/F}(x))$ for $x \geq 0$. We say that l is a lower ramification break for G if $G[u + \epsilon] \subsetneq G[u]$ for every $\epsilon > 0$. It is clear from the definitions that l is a lower ramification break if and only if $\phi_{E/F}(l)$ is an upper ramification break.

When $(K, A) \in \mathcal{B}$ the abelian subgroup A of $\text{Aut}_k(K)$ also has a ramification filtration. The lower ramification subgroups of A are defined by

$$A[x] = \{\sigma \in A : v_K(\sigma(T) - T) \geq x + 1\} \quad (2.2)$$

for $x \geq 0$. Since $A[x]$ has finite index in $A = A[0]$ for every $x \geq 0$, the function

$$\phi_A(x) = \int_0^x \frac{dt}{|A[0] : A[t]|} \quad (2.3)$$

is strictly increasing. Hence we can define the ramification subgroups of A with the upper numbering by $A(x) = A[\psi_A(x)]$, where $\psi_A(x) = \phi_A^{-1}(x)$. The upper and lower ramification breaks of A are defined in the same way as the upper and lower ramification breaks of $\text{Gal}(E/F)$. The lower ramification breaks of A are certainly integers, and Laubie's result (Corollary 1.2) together with the Hasse-Arf theorem imply that the upper ramification breaks of A are also integers.

For $E/F \in \mathcal{A}$ let $i(E/F)$ denote the smallest (upper or lower) ramification break of the extension E/F . The following basic result from ramification theory is presumably well-known (cf. [7, 3.2.5.5]).

Lemma 2.1 *Let $M/F \in \mathcal{A}$, let $L \in \mathcal{E}_{M/F}$, and let L'/L be a finite totally ramified abelian extension which is linearly disjoint from M/L . Assume that $M' = ML'$ has residue field k , so that $M'/F' \in \mathcal{A}$. Then $i(M'/F') \leq \psi_{F'/F}(i(M/F))$, with equality if the largest upper ramification break u of F'/F is less than $i(M/F)$.*

Proof: Set $G = \text{Gal}(M'/F)$, $H = \text{Gal}(M'/M)$, and $N = \text{Gal}(M'/F')$. Then $G = HN \cong H \times N$. Let $y = \phi_{F'/F}(i(M'/F'))$. Then

$$N = N(i(M'/F')) = N(\psi_{F'/F}(y)) = G(y) \cap N. \quad (2.4)$$

It follows that $G(y) \supset N$, and hence that $G/H = G(y)H/H = (G/H)(y)$. Therefore $y \leq i(M/F)$, which implies $i(M'/F') \leq \psi_{F'/F}(i(M/F))$.

If $u < i(M/F)$ then the group

$$(G/N)(i(M/F)) = G(i(M/F))N/N \quad (2.5)$$

is trivial. It follows that $G(i(M/F)) \subset N$, and hence that

$$N(\psi_{F'/F}(i(M/F))) = G(i(M/F)) \cap N = G(i(M/F)). \quad (2.6)$$

The restriction map from $\text{Gal}(M'/F') = N$ to $\text{Gal}(M/F) \cong G/H$ carries $G(i(M/F))$ onto

$$G(i(M/F))H/H = (G/H)(i(M/F)) = G/H. \quad (2.7)$$

Thus $N(\psi_{F'/F}(i(M/F))) = N$, so we have $i(M'/F') \geq \psi_{F'/F}(i(M/F))$. Combining this with the inequality proved above we get $i(M'/F') = \psi_{F'/F}(i(M/F))$. $\square$

Let $E/F \in \mathcal{A}$. Since E/F is an APF extension, the field of norms of E/F is defined: Let $\mathcal{E}_{E/F}$ denote the set of finite subextensions of E/F , and for $L', L \in \mathcal{E}_{E/F}$ such that $L' \supset L$ let $N_{L'/L} : L' \rightarrow L$ denote the norm map. The field of norms $X_F(E)$ of E/F is defined to be the inverse limit of $L \in \mathcal{E}_{E/F}$ with respect to the norms. We denote an element of $X_F(E)$ by $\alpha_{E/F} = (\alpha_L)_{L \in \mathcal{E}_{E/F}}$. Multiplication in $X_F(E)$ is defined componentwise, and addition is defined by the rule $\alpha_{E/F} + \beta_{E/F} = \gamma_{E/F}$, where

$$\gamma_L = \lim_{L' \in \mathcal{E}_{E/L}} N_{L'/L}(\alpha_{L'} + \beta_{L'}) \quad (2.8)$$

for $L \in \mathcal{E}_{E/F}$. We embed k into $X_F(E)$ as follows: Let F_0/F be the maximum tamely ramified subextension of E/F , and for $\zeta \in k$ let $\tilde{\zeta}_{F_0}$ be the Teichmüller lift of ζ in $\mathcal{O}_{F_0}$. Note that for any $L \in \mathcal{E}_{E/F_0}$ the degree of the extension L/F_0 is a power of p . Therefore there is a unique $\tilde{\zeta}_L \in L$ such that $\tilde{\zeta}_L^{[L:F_0]} = \tilde{\zeta}_{F_0}$. Define $f_{E/F}(\zeta)$ to be the unique element of $X_F(E)$ whose L component is $\tilde{\zeta}_L$ for every $L \in \mathcal{E}_{E/F_0}$. Then the map $f_{E/F} : k \rightarrow X_F(E)$ is a field embedding. By choosing a uniformizer for $X_F(E)$ we get a k -isomorphism $X_F(E) \cong k((T))$.

The ring of integers $\mathcal{O}_{X_F(E)}$ consists of those $\alpha_{E/F} \in X_F(E)$ such that $\alpha_L \in \mathcal{O}_L$ for all $L \in \mathcal{E}_{E/F}$ (or equivalently, for any $L \in \mathcal{E}_{E/F}$). A uniformizer $\pi_{E/F} = (\pi_L)_{L \in \mathcal{E}_{E/F}}$ for $X_F(E)$ gives a uniformizer π_L for each finite subextension L/F of E/F , and also a uniformizer $\pi_{M/F} = (\pi_L)_{L \in \mathcal{E}_{M/F}}$ of $X_F(M)$ for each infinite subextension M/F of E/F . The action of $\text{Gal}(E/F)$ on the fields $L \in \mathcal{E}_{E/F}$ induces a k -action of $\text{Gal}(E/F)$ on $X_F(E)$. By identifying $\text{Gal}(E/F)$ with the subgroup of $\text{Aut}_k(X_F(E))$ which it induces, we get the functor $\mathcal{F}(E/F) = (X_F(E), \text{Gal}(E/F))$ which was mentioned in (1.1).

Let E' be a finite extension of E such that $E'/F \in \mathcal{A}$. Then there is $M \in \mathcal{E}_{E/F}$ and a finite extension M' of M such that $E' = EM'$ and E, M' are linearly disjoint over M . We define an embedding $j : X_F(E) \rightarrow X_F(E')$ as follows. For $\alpha_{E/F} \in X_F(E)$ set $j(\alpha_{E/F}) = \beta_{E'/F}$, where $\beta_{E'/F}$ is the unique element of $X_F(E')$ such that $\beta_{LM'} = \alpha_L$ for all $L \in \mathcal{E}_{E/M}$ [7, 3.1.1]. The embedding j makes $X_F(E')$ into a finite separable extension of $X_F(E)$ of degree $[E' : E]$; in this setting we denote $X_F(E')$ by $X_{E/F}(E')$. If $E'' \supset E' \supset E$ are finite extensions such that $E''/F \in \mathcal{A}$ then $X_{E/F}(E'')/X_F(E)$

is a subextension of $X_{E/F}(E'')/X_F(E)$. Let D/E be an infinite extension such that $D/F \in \mathcal{A}$. Then $X_{E/F}(D)$ is defined to be the union of $X_{E/F}(E')$ as E' ranges over the finite subextensions of D/E .

For $E/F \in \mathcal{A}$ set $r(E/F) = \lceil \frac{p-1}{p} \cdot i(E/F) \rceil$. The proof of Theorem 1.1 depends on the following two results, the first of which was proved by Wintenberger:

Proposition 2.2 *Let $E/F \in \mathcal{A}$, let $L \in \mathcal{E}_{E/F_0}$, and define $\xi_L : \mathcal{O}_{X_F(E)} \rightarrow \mathcal{O}_L/\mathcal{M}_L^{r(E/L)}$ by $\xi_L(\alpha_{E/F}) = \alpha_L \pmod{\mathcal{M}_L^{r(E/L)}}$. Then*

- (a) ξ_L is a surjective ring homomorphism.
- (b) ξ_L induces the map $\zeta \mapsto \zeta^{p^{-a}}$ on k , where $a = v_p([L : F])$.

Proof: This is proved in Propositions 2.2.1 and 2.3.1 of [7]. □

Proposition 2.3 *Let $E/F \in \mathcal{A}$, let $L \in \mathcal{E}_{E/F}$, and let L'/L be a finite totally ramified abelian extension which is linearly disjoint from E/L . Assume that $E' = EL'$ has residue field k , so that $E'/L' \in \mathcal{A}$. Then the following diagram commutes, where the bottom horizontal map is induced by the inclusion $\mathcal{O}_L \hookrightarrow \mathcal{O}_{L'}$:*

$$\begin{array}{ccc} \mathcal{O}_{X_F(E)} & \xrightarrow{j} & \mathcal{O}_{X_{E/F}(E')} \\ \xi_L \downarrow & & \downarrow \xi_{L'} \\ \mathcal{O}_L/\mathcal{M}_L^{r(E/L)} & \longrightarrow & \mathcal{O}_{L'}/\mathcal{M}_{L'}^{r(E'/L')} \end{array} \quad (2.9)$$

Furthermore, for $\zeta \in k$ we have $j \circ f_{E/F}(\zeta) = f_{E'/F}(\zeta^{p^b})$, where $b = v_p([L' : L])$.

Proof: Using Lemma 2.1 we get

$$i(E'/L') \leq \psi_{L'/L}(i(E/L)) \leq [L' : L]i(E/L). \quad (2.10)$$

Thus $r(E'/L') \leq [L' : L]r(E/L)$, so the bottom horizontal map in the diagram is well-defined. Let $\alpha_{E/F} = (\alpha_M)_{M \in \mathcal{E}_{E/F}}$ be an element of $\mathcal{O}_{X_F(E)}$. Then $j(\alpha_{E/F})$ is the unique element of $\mathcal{O}_{X_{E/F}(E')}$ whose ML' -component is α_M for every $M \in \mathcal{E}_{E/L}$. In particular, the L' -component of $j(\alpha_{E/F})$ is α_L . Hence $\xi_L(\alpha_{E/F})$ and $\xi_{L'}(j(\alpha_{E/F}))$ are both congruent to α_L modulo $\mathcal{M}_{L'}^{r(E'/L')}$, which proves the commutativity of (2.9). The second claim follows from (2.9) and Proposition 2.2(b). □

3 Proof of Theorem 1.1

In this section we prove Theorem 1.1. To do this, we must show that the functor $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$ is essentially surjective and fully faithful.

We begin by showing that $\mathcal{F}$ is essentially surjective. Let A be a closed abelian subgroup of $\text{Aut}_k(K)$, where $K = k((T))$. Then A is a p -adic Lie group if and only if A is finitely generated. Since $\mathcal{F}$ induces an equivalence between the categories $\mathcal{A}_L$ and

$\mathcal{B}_L$, it suffices to prove that (K, A) lies in the essential image of $\mathcal{F}$ in the case where A is *not* finitely generated.

Let $F \cong k((T))$, let E/F be a finite totally ramified abelian extension, and let π be a uniformizer of E . Then for each $\sigma \in \text{Gal}(E/F)$ there is a unique $f_\sigma \in k[[T]]$ such that $\sigma(\pi) = f_\sigma(\pi)$. Let $a = v_p([E : F])$ and define

$$G(E/F, \pi) = \{\gamma \in \text{Aut}_k(K) : \gamma(T) = f_\sigma^{p^a}(T) \text{ for some } \sigma \in \text{Gal}(E/F)\}, \quad (3.1)$$

where $f_\sigma^{p^a}(T)$ is the power series obtained from $f_\sigma(T)$ by replacing the coefficients by their p^a powers. Then $G(E/F, \pi)$ is a subgroup of $\text{Aut}_k(K)$ which is isomorphic to $\text{Gal}(E/F)$.

Let $l_0 < l_1 < l_2 < \dots$ denote the positive lower ramification breaks of A . For $n \geq 0$ set $r_n = \lceil \frac{p-1}{p} \cdot l_n \rceil$ and let $\bar{\Gamma}_n$ denote the quotient of $\Gamma = \text{Aut}_k(K)$ by the lower ramification subgroup

$$\Gamma[r_n - 1] = \{\sigma \in \Gamma : \sigma(T) \equiv T \pmod{\mathcal{M}_K^{r_n}}\}. \quad (3.2)$$

Let $\mathcal{S}_n$ denote the set of pairs (E, π) such that

1. E/F is a totally ramified abelian subextension of F^{sep}/F such that $\text{Gal}(E/F)[l_n]$ is trivial. (Such an extension is necessarily finite.)
2. π is a uniformizer of E such that the image of $G(E/F, \pi)$ in $\bar{\Gamma}_n$ is equal to the image of A in $\bar{\Gamma}_n$.

Since there are only finitely many extensions E/F satisfying condition 1, and condition 2 depends only on the class of π modulo $\mathcal{M}_E^{r_n}$, the set $\mathcal{S}_n$ is compact. Using the following lemma we get a map from $\mathcal{S}_n$ to $\mathcal{S}_{n-1}$.

Lemma 3.1 *Let $n \geq 1$, let $(E, \pi) \in \mathcal{S}_n$, and let $\tilde{E}$ denote the fixed field of $\text{Gal}(E/F)[l_{n-1}]$. Then $(\tilde{E}, N_{E/\tilde{E}}(\pi)) \in \mathcal{S}_{n-1}$.*

Proof: It follows from the definitions that $\tilde{E}/F$ is a totally ramified abelian extension and that $\text{Gal}(\tilde{E}/F)[l_{n-1}]$ is trivial. Set $\tilde{\pi} = N_{E/\tilde{E}}(\pi)$, choose $\sigma \in \text{Gal}(E/F)$, and let $\tilde{\sigma}$ denote the restriction of σ to $\tilde{E}$. By [7, Prop. 2.2.1] the norm $N_{E/\tilde{E}}$ induces a ring homomorphism from $\mathcal{O}_E$ to $\mathcal{O}_{\tilde{E}}/\mathcal{M}_{\tilde{E}}^{r_{n-1}}$. Therefore

$$\tilde{\sigma}(\tilde{\pi}) = N_{E/\tilde{E}}(\sigma(\pi)) = N_{E/\tilde{E}}(f_\sigma(\pi)) \equiv f_\sigma^{p^b}(N_{E/\tilde{E}}(\pi)) \pmod{\mathcal{M}_{\tilde{E}}^{r_{n-1}}}, \quad (3.3)$$

where $b = v_p([E : \tilde{E}])$. Let $\tilde{a} = v_p([\tilde{E} : F])$ and let $\tilde{f}_\sigma \in k[[T]]$ be such that $\tilde{\sigma}(\tilde{\pi}) = \tilde{f}_\sigma(\tilde{\pi})$. Then by (3.3) we have

$$f_{\tilde{\sigma}}(T) \equiv f_\sigma^{p^b}(T) \pmod{T^{r_{n-1}}} \quad (3.4)$$

$$f_{\tilde{\sigma}}^{p^{\tilde{a}}}(T) \equiv f_\sigma^{p^a}(T) \pmod{T^{r_{n-1}}}. \quad (3.5)$$

It follows that $G(\tilde{E}/F, \tilde{\pi})$ and $G(E/F, \pi)$ have the same image in $\bar{\Gamma}_{n-1}$, and hence that $G(\tilde{E}/F, \tilde{\pi})$ and A have the same image in $\bar{\Gamma}_{n-1}$. $\square$

Since each $A/A[l_n]$ is finite there is a sequence $A_0 \leq A_1 \leq A_2 \leq \dots$ of finitely generated closed subgroups of A such that $A_n A[l_n] = A$ for all $n \geq 0$. Recall that $\mathcal{F}$ induces an equivalence of categories between $\mathcal{A}_L$ and $\mathcal{B}_L$. Since $(K, A_n) \in \mathcal{B}_L$, for $n \geq 0$ there exists $L_n/F_n \in \mathcal{A}_L$ such that $\mathcal{F}(L_n/F_n)$ is $\mathcal{B}$ -isomorphic to (K, A_n) . Since A_n is a normal subgroup of A , the action of A on K gives a $\mathcal{B}$ -action of A on the pair (K, A_n) . Since $\mathcal{F}(L_n/F_n) \cong (K, A_n)$ and $\mathcal{F}$ induces an equivalence between $\mathcal{A}_L$ and $\mathcal{B}_L$, this action is induced by a faithful $\mathcal{A}$ -action of A on L_n/F_n . Since $\text{Gal}(L_n/F_n) \cong A_n$ is finitely generated, and A is not finitely generated, this implies that $\text{Aut}_k(F_n)$ is not finitely generated. Hence F_n has characteristic p . Therefore we may assume $F_n = F$ and $L_n \subset F^{sep}$ with $F \cong k((T))$ fixed.

For $n \geq 0$ let

$$i_n : (K, A_n) \longrightarrow (X_F(L_n), \text{Gal}(L_n/F)) \quad (3.6)$$

be an $\mathcal{A}$ -isomorphism, and set $\pi_{L_n/F} = i_n(T)$. Let $E_n \subset L_n$ be the fixed field of $\text{Gal}(L_n/F)[l_n]$. Then either $E_n = L_n$, or $i(L_n/E_n) \geq l_n$, in which case $r(L_n/E_n) \geq r_n$. Therefore by Proposition 2.2(a) we have $(E_n, \pi_{E_n}) \in \mathcal{S}_n$, so $\mathcal{S}_n \neq \emptyset$. For $n \geq 1$ let $\nu_n : \mathcal{S}_n \rightarrow \mathcal{S}_{n-1}$ be the map defined by Lemma 3.1. Since each $\mathcal{S}_n$ is compact, by Tychonoff's theorem there exists a sequence of pairs $(E_n, \pi_{E_n}) \in \mathcal{S}_n$ such that $\nu_n(E_n, \pi_{E_n}) = (E_{n-1}, \pi_{E_{n-1}})$ for $n \geq 1$. It follows in particular that $F \subset E_0 \subset E_1 \subset E_2 \subset \dots$. Let $E_\infty = \bigcup_{n \geq 0} E_n$. Then E_∞ is a totally ramified abelian extension of F , and the uniformizers π_{E_n} for E_n induce a uniformizer $\pi_{E_\infty/F}$ for $X_F(E_\infty)$. Let τ denote the unique k -isomorphism from $K = k((T))$ to $X_F(E_\infty)$ such that $\tau(T) = \pi_{E_\infty/F}$. It follows from our construction that τ induces a $\mathcal{B}$ -isomorphism from (K, A) to

$$\mathcal{F}(E_\infty/F) = (X_F(E_\infty), \text{Gal}(E_\infty/F)). \quad (3.7)$$

Thus (K, A) lies in the essential image of $\mathcal{F}$, so $\mathcal{F}$ is essentially surjective.

We now show that $\mathcal{F}$ is faithful. Let E/F and E'/F' be elements of $\mathcal{A}$, and set $G = \text{Gal}(E/F)$ and $G' = \text{Gal}(E'/F')$. We need to show that the map

$$\Psi : \text{Hom}_{\mathcal{A}}(E/F, E'/F') \longrightarrow \text{Hom}_{\mathcal{B}}((X_F(E), G), (X_{F'}(E'), G')) \quad (3.8)$$

induced by the field of norms functor is one-to-one. Suppose $\rho_1, \rho_2 \in \text{Hom}_{\mathcal{A}}(E/F, E'/F')$ satisfy $\Psi(\rho_1) = \Psi(\rho_2)$. Let $\pi_{E/F} = (\pi_L)_{L \in \mathcal{E}_{E/F}}$ be a uniformizer for $X_F(E)$. Then $\Psi(\rho_1)(\pi_{E/F}) = \Psi(\rho_2)(\pi_{E/F})$, and hence $(\rho_1(\pi_L))_{L \in \mathcal{E}_{E/F}} = (\rho_2(\pi_L))_{L \in \mathcal{E}_{E/F}}$. It follows that $\rho_1(\pi_L) = \rho_2(\pi_L)$ for every $L \in \mathcal{E}_{E/F}$. Since ρ_1 and ρ_2 are k -algebra homomorphisms, this implies that $\rho_1 = \rho_2$.

It remains to show that $\mathcal{F}$ is full, i.e., that Ψ is onto. It follows from the arguments given in the proof of [6, Th. 2.1] that the codomain of Ψ is empty if $\text{char}(F) \neq \text{char}(F')$, and that Ψ is onto if G and G' are finitely generated. In particular, Ψ is onto if $\text{char}(F) = 0$ or $\text{char}(F') = 0$. If one of G, G' is finitely generated and the other is not then the codomain of Ψ is empty. Hence it suffices to prove that Ψ is onto in the case where $\text{char}(F) = \text{char}(F') = p$ and neither of G, G' is finitely generated.

We first show that every isomorphism lies in the image of Ψ . Let

$$\tau : (X_F(E), G) \longrightarrow (X_{F'}(E'), G') \quad (3.9)$$

be a $\mathcal{B}$ -isomorphism. For $n \geq 1$ let F_n denote the fixed field of $G[l_n] = G(u_n)$. If $\lim_{n \rightarrow \infty} l_n/[F_n : F] = \infty$ then an argument similar to that used in [5, §2] shows that τ is induced by an $\mathcal{A}$ -isomorphism from E/F to E'/F' . This limit condition holds for instance if $\text{char}(F) = p$ and $\text{Gal}(E/F)$ is finitely generated, but it can fail if $\text{Gal}(E/F)$ is not finitely generated. Therefore we use a different method to prove that τ lies in the image of Ψ , based on a characterization of F_n/F in terms of $(X_F(E), G)$.

Let d denote the F_n -valuation of the different of F_n/F , and let c be an integer such that $c > \phi_{F_n/F}(\frac{p}{p-1}(l_{n-1} + d))$. Since $G/G(c)$ is finite there exists a finitely generated closed subgroup H of G such that $HG(c) = G$. Let $M \subset E$ be the fixed field of H and set $M_n = F_n M$. Then F_n/F and M_n/M are finite abelian extensions. On the other hand, since G is not finitely generated, $\text{Gal}(M/F) \cong G/H$ is not finitely generated, and hence M/F is an infinite abelian extension.

Proposition 3.2 *Let $\pi_{E/F}$ be a uniformizer for $X_F(E)$, and recall that $\pi_{E/F}$ induces uniformizers π_F , π_{F_n} , $\pi_{M/F}$, and $\pi_{M_n/F}$ for F , F_n , $X_F(M)$, and $X_{M/F}(M_n)$. There exists a k -isomorphism $\zeta : X_{M/F}(M_n)/X_F(M) \rightarrow F_n/F$ such that*

1. $\zeta(\pi_{M/F}) = \pi_F$;
2. $\zeta(\pi_{M_n/F}) \equiv \pi_{F_n} \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}}$;
3. $\gamma \cdot \zeta(\pi_{M_n/F}) = \zeta(\gamma \cdot \pi_{M_n/F})$ for every $\gamma \in H$.

The proof of this proposition depends on the following lemma (cf. [1, p. 88]).

Lemma 3.3 *Let F be a local field, let $g(T) \in \mathcal{O}_F[T]$ be a separable monic Eisenstein polynomial, and let $\alpha \in F^{\text{sep}}$ be a root of $g(T)$. Set $E = F(\alpha)$ and let $d = v_E(g'(\alpha))$ be the E -valuation of the different of the extension E/F . Choose $\pi \in F^{\text{sep}}$ such that $v_E(g(\pi)) > d$. Then there is a root β for $g(X)$ such that $v_E(\pi - \beta) \geq v_E(g(\pi)) - d$.*

Proof: Let $\alpha_1, \alpha_2, \dots, \alpha_n$ be the roots of $g(T)$, and choose $1 \leq j \leq n$ to maximize $w = v_E(\pi - \alpha_j)$. For $1 \leq i \leq n$ we have

$$v_E(\pi - \alpha_i) \geq \min\{w, v_E(\alpha_j - \alpha_i)\}, \quad (3.10)$$

with equality if $w > v_E(\alpha_j - \alpha_i)$. Since $w \geq v_E(\pi - \alpha_i)$, this implies that for $i \neq j$ we have $v_E(\pi - \alpha_i) \leq v_E(\alpha_j - \alpha_i)$. Since

$$g(\pi) = (\pi - \alpha_1)(\pi - \alpha_2) \dots (\pi - \alpha_n), \quad (3.11)$$

we get

$$v_E(g(\pi)) \leq w + \sum_{\substack{1 \leq i \leq n \\ i \neq j}} v_E(\alpha_j - \alpha_i) = w + d. \quad (3.12)$$

Setting $\beta = \alpha_j$ gives $v_E(\pi - \beta) = w \geq v_E(g(\pi)) - d$. $\square$

Proof of Proposition 3.2: Since $HG(c) = G$ we have

$$i(M/F) \geq c > \phi_{F_n/F}(l_{n-1}) = u_{n-1}. \quad (3.13)$$

Therefore by Lemma 2.1 we get

$$i(M_n/F_n) = \psi_{F_n/F}(i(M/F)) \geq \psi_{F_n/F}(c), \quad (3.14)$$

and hence $r(M_n/F_n) \geq s$, where $s = \lceil \frac{p-1}{p} \cdot \psi_{F_n/F}(c) \rceil$. Let $g(T)$ be the minimum polynomial for $\pi_{M_n/F}$ over $X_F(M)$, and let $g_F(T) \in \mathcal{O}_F[T]$ be the polynomial obtained by applying the canonical map $\xi_F : X_F(M) \rightarrow F$ to the coefficients of $g(T)$. Since $g(\pi_{M_n/F}) = 0$, it follows from Propositions 2.2(a) and 2.3 that $v_{F_n}(g_F(\pi_{F_n})) \geq r(M_n/F_n) \geq s$. On the other hand, let $\mu : X_F(M) \rightarrow F$ be the unique k -algebra isomorphism such that $\mu(\pi_{M/F}) = \pi_F$. Then by Proposition 2.2(a) we have $\mu(\alpha_{M/F}) \equiv \alpha_F \pmod{\mathcal{M}_F^t}$ for all $\alpha_{M/F} \in \mathcal{O}_{X_F(M)}$, where $t = \lceil \frac{p-1}{p} \cdot c \rceil$. Let $g^\mu(T) \in \mathcal{O}_F[T]$ be the polynomial obtained by applying μ to the coefficients of $g(T)$. Then $g^\mu(T) \equiv g_F(T) \pmod{\mathcal{M}_F^t}$. Since $[F_n : F] \cdot c \geq \psi_{F_n/F}(c)$ we have $[F_n : F] \cdot t \geq s$; therefore since $v_{F_n}(g_F(\pi_{F_n})) \geq s$ we have $v_{F_n}(g^\mu(\pi_{F_n})) \geq s > l_{n-1} + d$. It follows from Lemma 3.3 that there is a root β of $g^\mu(T)$ such that $v_{F_n}(\pi_{F_n} - \beta) > l_{n-1}$. Therefore by Krasner's Lemma we have $F(\beta) \supset F(\pi_{F_n})$. Since

$$[F(\beta) : F] = \deg(g) = [F(\pi_{F_n}) : F] \quad (3.15)$$

we deduce that $F(\beta) = F(\pi_{F_n}) = F_n$. Since $\pi_{M_n/F}$ is a root of $g(T)$, and β is a root of $g^\mu(T)$, the isomorphism μ from $X_F(M)$ to F extends uniquely to an isomorphism ζ from $X_{M/F}(M_n)/X_F(M)$ to F_n/F such that $\zeta(\pi_{M_n/F}) = \beta \equiv \pi_{F_n} \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}}$.

We now show that ζ is H -equivariant. Let $\gamma \in H$ and define $\psi_\gamma \in k[[T]]$ by

$$\psi_\gamma(\pi_{M_n/F}) = \gamma \cdot \pi_{M_n/F} = (\gamma \cdot \pi_L)_{L \in \mathcal{E}_{M_n/F}}, \quad (3.16)$$

where we identify k with a subfield of $X_F(M)$ using the map $f_{M/F}$. Using Propositions 2.2 and 2.3 we get

$$\gamma \cdot \pi_{F_n} \equiv \psi_\gamma(\pi_{F_n}) \pmod{\mathcal{M}_{F_n}^{r(M_n/F_n)}}. \quad (3.17)$$

Since $\zeta(\pi_{M_n/F}) \equiv \pi_{F_n} \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}}$ and $r(M_n/F_n) \geq s \geq l_{n-1} + 1$ this implies

$$\zeta(\gamma \cdot \pi_{M_n/F}) = \zeta(\psi_\gamma(\pi_{M_n/F})) \quad (3.18)$$

$$= \psi_\gamma(\zeta(\pi_{M_n/F})) \quad (3.19)$$

$$\equiv \psi_\gamma(\pi_{F_n}) \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}} \quad (3.20)$$

$$\equiv \gamma \cdot \pi_{F_n} \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}} \quad (3.21)$$

$$\equiv \gamma \cdot \zeta(\pi_{M_n/F}) \pmod{\mathcal{M}_{F_n}^{l_{n-1}+1}}. \quad (3.22)$$

Since $\zeta(\gamma \cdot \pi_{M_n/F})$ and $\gamma \cdot \zeta(\pi_{M_n/F})$ are both roots of $g^\mu(T)$ we deduce that $\gamma \cdot \zeta(\pi_{M_n/F}) = \zeta(\gamma \cdot \pi_{M_n/F})$. Since ζ is k -linear and γ acts trivially on k , it follows that $\gamma \cdot \zeta(\alpha) = \zeta(\gamma \cdot \alpha)$ for all $\alpha \in X_{M/F}(M_n)$. $\square$

Since τ is an $\mathcal{A}$ -isomorphism, $\tau^* : G' \rightarrow G$ is a group isomorphism. For $\gamma \in G$ set $\gamma' = (\tau^*)^{-1}(\gamma)$, and for $N \leq G$ set $N' = (\tau^*)^{-1}(N)$. Then τ induces an isomorphism from $(X_F(E), N)$ to $(X_{F'}(E'), N')$. In particular, τ gives an isomorphism from $(X_F(E), H)$ to $(X_{F'}(E'), H')$. Using the isomorphism $X_{X_F(M)}(X_{M/F}(E)) \cong X_F(E)$ from [7, 3.4.1] we get an isomorphism

$$\tau_H : (X_{X_F(M)}(X_{M/F}(E)), H) \longrightarrow (X_{X_{F'}(M')}(X_{M'/F'}(E')), H'), \quad (3.23)$$

where $M' \subset E'$ is the fixed field of H' . Since H is an abelian p -adic Lie group, it follows from [2, 5, 6] that τ_H is induced by an $\mathcal{A}$ -isomorphism

$$\rho : X_{M/F}(E)/X_F(M) \longrightarrow X_{M'/F'}(E')/X_{F'}(M'). \quad (3.24)$$

By restricting ρ we get an isomorphism

$$\tilde{\rho} : X_{M/F}(M_n)/X_F(M) \longrightarrow X_{M'/F'}(M'_n)/X_{F'}(M'), \quad (3.25)$$

where $M'_n = (M')_n = F'_n M'$ is the fixed field of $H'[l_n] = H[l_n]'$. Furthermore, for $\gamma \in H$ and $\alpha \in X_{M/F}(M_n)$ we have $\tilde{\rho}(\gamma(\alpha)) = \gamma'(\tilde{\rho}(\alpha))$.

Let $\pi_{E/F}$ be a uniformizer for $X_F(E)$, set $\pi_{E'/F'} = \tau(\pi_{E/F})$, and let

$$\zeta : X_{M/F}(M_n)/X_F(M) \longrightarrow F_n/F \quad (3.26)$$

$$\zeta' : X_{M'/F'}(M'_n)/X_{F'}(M') \longrightarrow F'_n/F' \quad (3.27)$$

be the isomorphisms given by Proposition 3.2. Then $\omega_n = \zeta' \circ \tilde{\rho} \circ \zeta^{-1}$ gives a k -linear isomorphism from F_n/F to F'_n/F' . It follows from Proposition 3.2 that

$$\omega_n(\pi_{F_n}) \equiv \pi_{F'_n} \pmod{\mathcal{M}_{F'_n}^{l_{n-1}+1}}, \quad (3.28)$$

and that for all $\gamma \in H$ we have

$$\omega_n(\gamma(\pi_{F_n})) = \gamma'(\omega_n(\pi_{F_n})). \quad (3.29)$$

Since the restriction map from $H = \text{Gal}(E/M)$ to $\text{Gal}(F_n/F)$ is onto, (3.29) is actually valid for all $\gamma \in G$.

Let $\mathcal{T}_n$ denote the set of k -isomorphisms $\omega_n : F_n/F \rightarrow F'_n/F'$ which satisfy (3.28) and (3.29) for all $\gamma \in G$. Since l_{n-1} is the only ramification break of F'_n/F'_{n-1} we have $\psi_{F'_n/F'_{n-1}}(l_{n-1}) = l_{n-1}$. Therefore by (3.28) and [4, V §6, Prop. 8], for any $\omega_n \in \mathcal{T}_n$ we have

$$N_{L'_n/L'_{n-1}}(\omega_n(\pi_{F_n})) \equiv N_{L'_n/L'_{n-1}}(\pi_{F'_n}) \pmod{\mathcal{M}_{F'_{n-1}}^{l_{n-1}+1}}. \quad (3.30)$$

Since $N_{L_n/L_{n-1}}(\pi_{F_n}) = \pi_{F_{n-1}}$ and $N_{L'_n/L'_{n-1}}(\pi_{F'_n}) = \pi_{F'_{n-1}}$, it follows from (3.30) and (3.29) that

$$\omega_n(\pi_{F_{n-1}}) \equiv \pi_{F'_{n-1}} \pmod{\mathcal{M}_{F'_{n-1}}^{l_{n-1}+1}}. \quad (3.31)$$

Therefore restriction induces a map from $\mathcal{T}_n$ to $\mathcal{T}_{n-1}$.

Define a metric on $\mathcal{T}_n$ by setting $d(\omega_n, \tilde{\omega}_n) = 2^{-a}$, where $a = v_{F'_n}(\omega_n(\pi_{F_n}) - \tilde{\omega}_n(\pi_{F_n}))$. Then $\mathcal{T}_n$ is easily seen to be compact, and we showed above that $\mathcal{T}_n$ is nonempty. Therefore by Tychonoff's theorem there is a sequence $(\omega_n)_{n \geq 1}$ with $\omega_n \in \mathcal{T}_n$ and $\omega_n|_{F_{n-1}} = \omega_{n-1}$. Since $E = \bigcup_{n \geq 1} F_n$ and $E' = \bigcup_{n \geq 1} F'_n$ the isomorphisms $\omega_n : F_n/F \rightarrow F'_n/F'$ combine to give an $\mathcal{A}$ -isomorphism $\Omega : E/F \rightarrow E'/F'$. Let $\theta = \Psi(\Omega)$ be the $\mathcal{B}$ -isomorphism induced by Ω and let $m_n = \min\{l_{n-1} + 1, r(E/F_n)\}$. It follows from (3.28) and Proposition 2.2(a) that

$$\theta(\pi_{E/F}) \equiv \pi_{E'/F'} \pmod{\mathcal{M}_{X_{F'}(E')}^{m_n}} \quad (3.32)$$

for every $n \geq 1$. Since $\lim_{n \rightarrow \infty} m_n = \infty$ we get $\theta(\pi_{E/F}) = \pi_{E'/F'} = \tau(\pi_{E/F})$. Hence $\tau = \theta = \Psi(\Omega)$.

Now let σ be an arbitrary element of $\text{Hom}_{\mathcal{B}}((X_F(E), G), (X_{F'}(E'), G'))$. Since $X_{F'}(E')$ is a finite separable extension of $\sigma(X_F(E))$, by [7, 3.2.2] there is a finite separable extension $\tilde{E}'$ of E such that σ extends to an isomorphism

$$\tau : X_{E/F}(\tilde{E}') \longrightarrow X_{F'}(E'). \quad (3.33)$$

It follows that each $\gamma' \in G'$ induces an automorphism $\tilde{\gamma}$ of $X_{E/F}(\tilde{E}')$ whose restriction to $X_F(E)$ is $\sigma^*(\gamma') \in G$. Since $X_{E/F}(F^{sep})$ is a separable closure of $X_F(E)$ [7, Cor. 3.2.3], $\tilde{\gamma}$ can be extended to an automorphism $\bar{\gamma}$ of $X_{E/F}(F^{sep})$. Since $\bar{\gamma}$ stabilizes $X_F(E)$, and $\bar{\gamma}|_{X_F(E)} = \sigma^*(\gamma')$ is induced by an element of $G = \text{Gal}(E/F)$, it follows from [7, Rem. 3.2.4] that $\bar{\gamma}$ is induced by an element of $\text{Gal}(F^{sep}/F)$, which we also denote by $\bar{\gamma}$. Since $\bar{\gamma}$ stabilizes $X_{E/F}(\tilde{E}')$, it stabilizes $\tilde{E}'$ as well. Thus $\bar{\gamma}|_{\tilde{E}'}$ is an element of $\text{Aut}_k(\tilde{E}')$ which is uniquely determined by γ' . Since $\bar{\gamma}|_{\tilde{E}'}$ induces the automorphism $\tilde{\gamma}$ of $X_{E/F}(\tilde{E}')$, we denote $\bar{\gamma}|_{\tilde{E}'}$ by $\tilde{\gamma}$ as well.

Let $\tilde{F}'$ denote the subfield of $\tilde{E}'$ which is fixed by the group $\tilde{G}' = \{\tilde{\gamma} : \gamma' \in G'\}$. Then $\tilde{F}' \supset F$, so $\tilde{E}'/\tilde{F}'$ is a Galois extension. Since the image of $\tilde{G}' \cong G'$ in G is open, and $\tilde{E}'/E$ is a finite extension, it follows that $\tilde{F}'$ is a finite separable extension of F , and $\tilde{G}' = \text{Gal}(\tilde{E}'/\tilde{F}')$. In particular, $\tilde{F}' \cong k((T))$ is a local field with residue field k . Hence $(X_{\tilde{F}'}(\tilde{E}'), \tilde{G}')$ is an object in $\mathcal{B}$, and τ gives a $\mathcal{B}$ -isomorphism from $(X_{\tilde{F}'}(\tilde{E}'), \tilde{G}')$ to $(X_{F'}(E'), G')$. By the arguments given above we see that τ is induced by an $\mathcal{A}$ -isomorphism $\Omega : \tilde{E}'/\tilde{F}' \rightarrow E'/F'$. Furthermore, the embedding $E \hookrightarrow \tilde{E}'$ induces an $\mathcal{A}$ -morphism $i : E/F \rightarrow \tilde{E}'/\tilde{F}'$. Let

$$\alpha : (X_F(E), G) \longrightarrow (X_{\tilde{F}'}(\tilde{E}'), \tilde{G}') \quad (3.34)$$

be the $\mathcal{B}$ -morphism induced by i . Then $\sigma = \tau \circ \alpha = \Psi(\Omega \circ i)$.

References

- [1] K. Keating, Extensions of local fields and truncated power series, *J. Number Theory* **116** (2006), 69–101.
- [2] F. Laubie, Extensions de Lie et groupes d'automorphismes de corps locaux, *Compositio Math.* **67** (1988), 165–189.

- [3] F. Laubie, Ramification des groupes abéliens d’automorphismes de $\mathbb{F}_q((X))$, *Canad. Math. Bull.* **50** (2007), 594–597.
- [4] J.-P. Serre, *Corps Locaux*, Hermann, Paris (1962).
- [5] J.-P. Wintenberger, Automorphismes des corps locaux de caractéristique p , *J. Théor. Nombres Bordeaux* **16** (2004), 429–456.
- [6] J.-P. Wintenberger, Extensions abéliennes et groupes d’automorphismes de corps locaux, *C. R. Acad. Sci. Paris Sér. A-B* **290** (1980), A201–A203.
- [7] J.-P. Wintenberger, Le corps des normes de certaines extensions infinies de corps locaux; applications, *Ann. Sci. École Norm. Sup. (4)* **16** (1983), 59–89.